

## University of Groningen

### **Draft genome sequence of *Bacillus mycoides* M2E15, a strain isolated from the endosphere of potato**

Yi, Yanglei; de Jong, Anne; Spoelder, Jan; Elzenga, J Theo M; van Elsas, Jan Dirk; Kuipers, Oscar P

*Published in:*  
Genome Announcements

*DOI:*  
[10.1128/genomeA.00031-16](https://doi.org/10.1128/genomeA.00031-16)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2016

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Yi, Y., de Jong, A., Spoelder, J., Elzenga, J. T. M., van Elsas, J. D., & Kuipers, O. P. (2016). Draft genome sequence of *Bacillus mycoides* M2E15, a strain isolated from the endosphere of potato. *Genome Announcements*, 4(1), 1-2. [e00031-16]. <https://doi.org/10.1128/genomeA.00031-16>

#### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

#### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

# Draft Genome Sequence of *Bacillus mycoides* M2E15, a Strain Isolated from the Endosphere of Potato

Yanglei Yi,<sup>a</sup> Anne de Jong,<sup>a</sup> Jan Spoelder,<sup>b,d</sup> J. Theo M. Elzenga,<sup>b</sup> Jan Dirk van Elsas,<sup>c</sup> Oscar P. Kuipers<sup>a</sup>

Molecular Genetics, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, Groningen, the Netherlands<sup>a</sup>; Plant Physiology, Groningen Institute for Evolutionary Life Sciences, University of Groningen, Groningen, the Netherlands<sup>b</sup>; Microbial Ecology, Groningen Institute for Evolutionary Life Sciences, University of Groningen, Groningen, the Netherlands<sup>c</sup>; HLB Research & Consultancy in Agriculture, Wijster, the Netherlands<sup>d</sup>

**We present the draft genome sequence of *Bacillus mycoides* M2E15, a bacterium isolated from potato endosphere. Analysis of the 6.08-Mbp draft genome sequence identified 6,386 protein-encoding sequences, including potential plant growth promoting genes. Specifically, genes for proteins involved in phosphate utilization, iron acquisition, and bacteriocin production were identified.**

Received 7 January 2016 Accepted 11 January 2016 Published 25 February 2016

**Citation** Yi Y, de Jong A, Spoelder J, Elzenga JTM, van Elsas JD, Kuipers OP. 2016. Draft genome sequence of *Bacillus mycoides* M2E15, a strain isolated from the endosphere of potato. *Genome Announc* 4(1):e00031-16. doi:10.1128/genomeA.00031-16.

**Copyright** © 2016 Yi et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Oscar P. Kuipers, o.p.kuipers@rug.nl.

*Bacillus mycoides* is a rod-shaped chain-forming bacterium which is associated with the *Bacillus cereus* group. On agar plates, it forms unique rhizoid colonies resulting from cells that are linked end-to-end. Several reports on *B. mycoides* have described its plant-growth-promoting effects on sugar beet, cucurbits, and tobacco (1–3). We found *B. mycoides* to be present in the endosphere of potato without causing visible signs of disease. In order to understand the genetic make-up of this bacterium, we present the draft genome of the endophytic strain M2E15, which was isolated from the roots of potato (cultivar Miranda) in Wijster, the Netherlands.

Strain M2E15 was grown overnight in Luria-Bertani (LB) broth at 30°C, 200 rpm. Bacterial cells were harvested at the exponential growth phase and lysed with lysozyme. After RNase treatment, proteins were removed by proteinase K digestion. DNA was extracted from the lysate by phenol-chloroform treatment and recovered by isopropanol precipitation. Purified genomic DNA was sequenced using the MiSeq sequencing system of Illumina (4), yielding 250 bp paired-end reads with a mean library size of 400 bp. *De novo* assembly was performed using Velvet (4). Prediction of protein-encoding regions and automatic functional annotation was performed using the Rapid Annotations using Subsystem Technology (RAST) server (5). Moreover, the bacteriocin identification tool Bagel3 was used (6). The assembled genome of strain M2E15 consisted of 236 contigs with a total size of 6,077,838 bp. The chromosome harbors a total of 6,386 putative coding sequences (CDS) and 37 genes for tRNAs.

The *B. mycoides* M2E15 genome presents several genes that may be related to the solubilization of phosphate in soil. These include glucose dehydrogenase, lactate dehydrogenase, and citrate synthase, which are all involved in the production of organic acids. Organic acid production and release is considered to be a key mechanism for mineral phosphate solubilization. Moreover, a purple acid phosphatase gene, alkaline phosphatase and three 5'-nucleotidase genes were found, which may aid in the mineraliza-

tion of organic phosphorus compounds (7). Several genes encoding proteins for the biosynthesis of exopolysaccharides were found in strain M2E15; these have considerable similarity with the *B. subtilis* eps operon, which is involved in biofilm formation, potentially aiding the organism in its interaction with plant roots (8). Genes for plant cell wall degradation enzymes, such as endoglucanase and glycoside hydrolase, are also present, suggesting its ability to degrade plant polymers (9, 10). Furthermore, the M2E15 genome was found to contain eleven chitinase genes, which may play a role in from the degradation of fungal and/or insect polymers (11). Finally, four bacteriocin gene clusters were predicted by BAGEL 3 (6).

**Nucleotide sequence accession numbers.** This whole-genome shotgun project has been deposited at GenBank under the accession number [LLWA00000000](https://www.ncbi.nlm.nih.gov/nuclseq/LLWA00000000/). The version described in this paper is version LLWA01000000.

## ACKNOWLEDGMENTS

This work was supported by a scholarship to Y.Y. from the China Scholarship Council (201306300040).

## REFERENCES

- Fravel DR, Spurr H, Harvey J. 1977. Biocontrol of tobacco brown-spot disease by *Bacillus cereus* subsp. *mycoides* in a controlled environment. *Phytopathology* 77:930–932. [http://dx.doi.org/10.1094/Phyto-67-930](https://doi.org/10.1094/Phyto-67-930).
- Bargabus R, Zidack N, Sherwood J, Jacobsen B. 2002. Characterisation of systemic resistance in sugar beet elicited by a non-pathogenic, phyllosphere-colonizing *Bacillus mycoides*, biological control agent. *Physiol Mol Plant* P61:289–298.
- Neher OT, Johnston MR, Zidack NK, Jacobsen BJ. 2009. Evaluation of *Bacillus mycoides* isolate BmJ and *B. mojavensis* isolate 203-7 for the control of anthracnose of cucurbits caused by *Glomerella cingulata* var. *orbiculare*. *Biol Contr* 48:140–146. [http://dx.doi.org/10.1016/j.biocontrol.2008.08.012](https://doi.org/10.1016/j.biocontrol.2008.08.012).
- Zerbino DR, Birney E. 2008. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. *Genome Res* 18:821–829. [http://dx.doi.org/10.1101/gr.074492.107](https://doi.org/10.1101/gr.074492.107).

5. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
6. van Heel AJ, de Jong A, Montalbán-López M, Kok J, Kuipers OP. 2013. BAGEL3: automated identification of genes encoding bacteriocins and (non-) bactericidal posttranslationally modified peptides. *Nucleic Acids Res* 41:W448–W453. <http://dx.doi.org/10.1093/nar/gkt391>.
7. Rodríguez H, Fraga R. 1999. Phosphate solubilizing bacteria and their role in plant growth promotion. *Biotechnol Adv* 17:319–339. [http://dx.doi.org/10.1016/S0734-9750\(99\)00014-2](http://dx.doi.org/10.1016/S0734-9750(99)00014-2).
8. Meneses CH, Rouws LF, Simões-Araújo JL, Vidal MS, Baldani JJ. 2011. Exopolysaccharide production is required for biofilm formation and plant colonization by the nitrogen-fixing endophyte *Gluconacetobacter diazotrophicus*. *Mol Plant Microbe* 24:1448–1458. <http://dx.doi.org/10.1094/MPMI-05-11-0127>.
9. Pedrosa FO, Monteiro RA, Wassem R, Cruz LM, Ayub RA, Colauto NB, Fernandez MA, Fungaro MHP, Grisard EC, Hungria M, Madeira HM, Nodari RO, Osaku CA, Petzl-Erler ML, Terenzi H, Vieira LG, Steffens MB, Weiss VA, Pereira LF, Almeida MI. 2011. Genome of *Herbaspirillum seropedicae* strain SmR1, a specialized diazotrophic endophyte of tropical grasses. *PLoS Genet* 7:e1002064. <http://dx.doi.org/10.1371/journal.pgen.1002064>.
10. Krause A, Ramakumar A, Bartels D, Battistoni F, Bekel T, Boch J, Böhm M, Friedrich F, Hurek T, Krause L, Linke B, McHardy AC, Sarkar A, Schneiker S, Syed AA, Thauer R, Vorhölter FJ, Weidner S, Pühler A, Reinhold-Hurek B, Kaiser O, Goesmann A. 2006. Complete genome of the mutualistic, N<sub>2</sub>-fixing grass endophyte *Azoarcus* sp. strain BH72. *Nat Biotechnol* 24:1385–1391. <http://dx.doi.org/10.1038/nbt1243>.
11. Krechel A, Faupel A, Hallmann J, Ulrich A, Berg G. 2002. Potato-associated bacteria and their antagonistic potential towards plant-pathogenic fungi and the plant-parasitic nematode *Meloidogyne incognita* (Kofoid & White) Chitwood. *Can J Microbiol* 48:772–786. <http://dx.doi.org/10.1139/w02-071>.